

Estimate of the total kinetic power and age of extragalactic jet by its cocoon dynamics: The case of Cygnus A

M. Kino and N. Kawakatu

SISSA, via Beirut 2-4, 34014 Trieste, Italy

submitted to MNRAS — — —

ABSTRACT

We examine the constraints imposed on the total kinetic power (L_j) and the age (t_{age}) of relativistic jets in FR II radio sources in a new way. We solve the dynamical expansion of its cocoon embedded in the intra-cluster medium (ICM) and obtain the analytic solution of its physical quantities in terms of L_j and t_{age} . The estimate of L_j and t_{age} is done by the comparison of the model and the observed shape of the cocoon. The analysis is focused on Cygnus A and we find that (i) the source age is $3 \text{ Myr} < t_{\text{age}} < 30 \text{ Myr}$ and (ii) the total kinetic power of the jet is estimated as $0.2 \times 10^{46} \text{ erg s}^{-1} < L_j < 1 \times 10^{48} \text{ erg s}^{-1}$ which is larger than 1% of the Eddington luminosity of Cygnus A.

Key words: plasmas — radiation mechanisms: non-thermal — galaxies: active — galaxies: individual: Cygnus A

1 INTRODUCTION

After the detections of inverse Compton component in X -ray from FR II radio sources, our knowledge of energetics, especially on the kinetic power of non-thermal electrons, is progressed in recent years (e.g., Leahy and Gizani 2001; Harris and Krawczynski 2002; Hardcastle et al. 2002; Kataoka et al. 2003; Croston et al. 2005). When we explore further physical conditions in the jets, we encounter a big difficulty to constrain on the fraction such as thermal electrons and/or protons co-existing with non-thermal electrons because it is hard to observe these component (e.g., Celotti et al. 1998; Sikora and Madejski 2002). This problem prevented us from estimating the *total* mass and energy flux ejected from a central engine. To conquer this, a simple procedure is proposed in the study of jets in FR II sources in Kino and Takahara (2004) (hereafter KT04). In the pressure and mass density of jet, the contributions from the invisible components are also involved. Hence the rest mass density and energy density estimated from the shock dynamics definitely prove the quantities of total plasma. Also for radio bubbles in cluster cores, the similar dynamical approach has been adopted to constrain on their physical state (Fabian et al. 2002; Dunn and Fabian 2004).

In this paper, we will explore the cocoon dynamics in FR II radio sources as a robust tool to know the total kinetic power L_j and source age t_{age} . In §2, we discuss our analytical solution of cocoon expansion which includes the effect of radial dependence of the rest mass density of the surrounding Intra Cluster Medium (ICM). In §3, by com-

paring the analytic solution and observation of the cocoon, we explore the kinetic power of Cygnus A, which is one of the best-studied FR II source. Conclusions are given in §4.

2 DYNAMICS OF COCOON

We deal with the cocoon dynamics in FR II radio source. The adopted basic equations are almost the same as those in Begelman and Cioffi (1989) (hereafter BC89). The main differences between BC89 and the present work are (i) we explicitly solve the physical quantities as a function of L_j , and t_{age} , (ii) we take account of the effect of radial profile of the mass density of ICM, and, as a result, (iii) the growth of the cocoon head A_h is consistently solved from the basic equations.

2.1 Basic Assumptions

Our main assumptions are as follows; (1) We limit our attention a jet relativistic speed ($v_j = c$), (2) The jet supplies a constant L_j in time, (3) We focus on the over-pressured cocoon phase, and (4) We assume that the magnetic fields are passive and ignore their dynamical effect. For (1), although it is still under debate, some jets are suggested to be relativistic (e.g., Tavecchio et al. 2000; Celotti, Ghisellini, & Chianberge 2001). Since little is known about the evolution of L_j , the assumption (2) is adopted as a first-step working hypothesis. The assumption of (3) is automatically guaranteed by the sideways expansion of the cocoon (e.g., Cioffi

and Blondin 1992). The assumption of (4) is based on the results that a multi-frequency analysis of radio galaxies show that the energy density of magnetic field tend to be smaller than that of non-thermal electrons (e.g., Leahy and Giani 2001; Isobe et al. 2002).

2.2 Basic Equations

Model parameters are L_j and t_{age} . Unknown physical quantities are v_h , v_c , P_c , and A_h (or A_c) which are the advance velocity of the cocoon head, the velocity of cocoon sideways expansion, the pressure of the cocoon, the cross sectional area of the head part of the cocoon (or the cross sectional area of cocoon body), respectively (see Fig. 1). Equation of motion along the jet axis, and sideways expansion, and energy conservation in the cocoon are given by

$$\frac{L_j}{v_j} = \rho_a(r_h) v_h^2(t) A_h(t), \quad (1)$$

$$P_c(t) = \rho_a(r_c) v_c(t)^2, \quad (2)$$

$$\frac{P_c(t) V_c(t)}{\hat{\gamma}_c - 1} \simeq 2 L_j t, \quad (3)$$

where $r_h(t) = \int_{t_{\min}}^t v_h(t') dt'$, $r_c(t) = \int_{t_{\min}}^t v_c(t') dt'$, $V_c(t) = 2 \int_{t_{\min}}^t A_c(t') v_h(t') dt'$, $t_{\min}$, and $\hat{\gamma}_c$ are the length from the center of the galaxy to the head of the cocoon, the radius of the cocoon body, the volume of the cocoon, the start time of source evolution and specific heat ratio of the plasma inside the cocoon, respectively. The declining mass density of ICM ρ_a is given by $\rho_a(r) = \bar{\rho}_a(r/r_0)^{-\alpha}$ where r_0 and $\bar{\rho}_a$ are reference position and the ICM mass density at r_0 , respectively. We set r_0 as $r_h(t_{\text{age}})$ in throughout this paper. Most of the kinetic energy is deposited in the cocoon, which is initially suggested by Scheuer (1974) and recent studies of hot spots also shows the radiative efficiency is very small (e.g., KT04). Following to Cioffi and Blondin (1992), we add the factor of $1/(\hat{\gamma}_c - 1)$ in Eq. (3) to express the amount of the deposited internal energy. In other words, this corresponds to the neglect of the PdV work because of its smallness.

The numbers of quantities are 4, while those of basic Eqs. are 3. Hence, we set $A_c(t) \propto t^X$ where X as a free parameter. We can constrain on the value of X from observations. A specific case is shown in §3. Hence, we obtain v_h , v_c , P_c , and A_h by using a free parameter X .

As a subsidiary equation, the area of the cocoon body is given by $A_c(t) = \pi \left(\int_{t_{\min}}^t v_c(t') dt' \right)^2$. It is clear that the change of the unknown from A_c to A_h does not change the result. However note that when we choose A_h as an unknown instead, we cannot obtain the solution for $\alpha = 2$.

2.3 Analytic Solution

We assume that physical quantities have a power law dependence in time and the coefficient of each physical quantity is barred quantity. Each quantity has the form of $A = \bar{A} (t/t_{\text{age}})^Y$ where Y is an arbitrary index. The time evolution of v_c is

$$v_c(t) = \bar{v}_c \left(\frac{t}{t_{\text{age}}} \right)^{0.5X-1} = \frac{\bar{A}_c^{1/2}}{t_{\text{age}}} \mathcal{C}_{vc} \left(\frac{t}{t_{\text{age}}} \right)^{0.5X-1}, \quad (4)$$

for a given A_c . With this, the analytic form of cocoon quantities in decreasing ICM density is obtained as follows;

$$P_c(t) = \bar{\rho}_a \bar{v}_c^2 \mathcal{C}_{pc} \left(\frac{\bar{v}_c}{v_0} \right)^{-\alpha} \left(\frac{t}{t_{\text{age}}} \right)^{X(1-\alpha/2)-2}, \quad (5)$$

$$v_h(L_j, t) = \frac{L_j}{\bar{\rho}_a \bar{v}_c^2 A_c} \mathcal{C}_{vh} \left(\frac{\bar{v}_c}{v_0} \right)^{\alpha} \left(\frac{t}{t_{\text{age}}} \right)^{X(-2+0.5\alpha)+2}, \quad (6)$$

$$A_h(L_j, t) = \frac{L_j}{v_j \bar{\rho}_a \bar{v}_h^2 A_c} \mathcal{C}_{ah} \left(\frac{\bar{v}_h}{v_0} \right)^{\alpha} \left(\frac{t}{t_{\text{age}}} \right)^{X(\alpha-2)(-2+0.5\alpha)+3\alpha-4}, \quad (7)$$

where $\mathcal{C}_{vh} = (\hat{\gamma}_c - 1)[3 - (1 - 0.5\alpha)X](0.5X)^{-\alpha}$, $\mathcal{C}_{vc} = 0.5X/\pi^{1/2}$, $\mathcal{C}_{pc} = (0.5X)^{\alpha}$, and $\mathcal{C}_{ah} = [X(-2+0.5\alpha)+3]^{-\alpha}$, respectively. $v_0 \equiv r_h/t_{\text{age}}$ corresponds to the head speed for constant velocity in time.

Here we use the conditions of $0.5X > 0$ and $X(-2+0.5\alpha)+3 > 0$ which make the contribution at $t_{\min}$ in the integrations of r_h and r_c small enough. The case we focus on in §3 is that $X = 12/7$ and $\alpha = 1.5$, which clearly satisfies these conditions.

First, let us consider the evolution of cocoon. The growth of both A_h and A_c must be positive. As for the cocoon expansion speeds, three different behaviors are theoretically predicted such as (I) *accelerated-head* ($X(-2+0.5\alpha)+2 > 0$), (II) *constant-head* ($X(-2+0.5\alpha)+2 = 0$), and (III) *decelerated-head* ($X(-2+0.5\alpha)+2 < 0$). The case of (I), (II), and (III) correspond to $X < 1$, $X = 1$, and $X > 1$ for $\alpha = 0$, while in the case of $\alpha = 2$ (I), (II), and (III) correspond to $X < 2$, $X = 2$, and $X > 2$, respectively. Related to this, it is useful to express the aspect ratio of the cocoon $\frac{r_c}{r_h} \equiv \mathcal{R}$ as a function of time. This is written as

$$\mathcal{R}(t) = \frac{X(-2+0.5\alpha)+3}{0.5X} \frac{\bar{v}_c}{\bar{v}_h} \left(\frac{t}{t_{\text{age}}} \right)^{X(2.5-0.5\alpha)-3}. \quad (8)$$

It is worth to note that the solution describes not only the self-similar evolution (e.g., Kaiser and Alexander 1997; Bicknell, Dopita and O'Dea 1997) but also the non self-similar one. Although it is not concerned with in this paper, the evolution of $\mathcal{R}(t)$ may probe the evolution of radio galaxies such as ‘‘compact symmetric objects (CSO)’’ which are thought to be the progenitors of the FR II source (e.g., Fanti et al. 1996; Readhead et al. 1996a). Note that, observationally, the large deviation from $\mathcal{R} \sim \mathcal{O}(1)$ does not seem to be unnatural for actual radio sources (e.g., Readhead et al. 1996b).

Next, let us consider the α and X dependences on the quantities based on Eqs. (1), (2) and (3). Here we fix the physical quantities at $t = t_{\text{age}}$. With regard to the α dependence in fixed X , larger α leads to larger ρ_a , slower v_h , same v_c , larger P_c , during $t < t_{\text{age}}$. We can understand it as follows. Larger α leads to stronger the deceleration effect on the head speed v_h due to larger ρ_a . Larger α also predict larger P_c in order to keep the same velocity of sideways expansion v_c . In Fig. 2, we show the effect of varying α when fixing $X = 1$. It shows that larger α requires larger L_j in order to plow the larger amount of ICM.

Next we consider the X dependence. In fixed α , larger X lead to faster v_h , slower v_c , smaller P_c , and smaller A_h during $t < t_{\text{age}}$. We can explain it as follows. Larger X leads to slower sideways expansion v_c and smaller A_c by definition.

From the equation of motion to the sideways expansion, it is clear that smaller P_c is required. To satisfy the energy equation at the same time, the faster velocity of v_h is needed. This is realized by smaller area head of the cocoon A_h .

Observationally, little is argued about the emission from the cocoon itself. Recently, Readhead et al. (1996b) report the cocoon emission from the CSO 2352+495. It is visible in the 610 MHz image and the aspect-ratio is about $\mathcal{R}(t_{\min}) \sim 1/2$. Because of the lack of our knowledge of the cocoon emissions, here we examine the case of $\mathcal{R}(t_{\min}) = 1, 1/2$, and $1/4$ for the moment. In Fig. 3, we show the effect of varying X with fixing $\alpha = 1.5$ and $\mathcal{R}(t_{\text{age}}) = 1/2$. The case of $X = 1.62, 12/7$, and 1.80 correspond to the $\mathcal{R}(t_{\min}) = 1, 1/2$, and $1/4$ respectively. We can verify the shift of L_j to smaller range as the X increases.

Finally, we compare our solution with the previous works. The solution for flat ICM density by BC89 model corresponds to the case of $X = 1$ and $\alpha = 0$ and the quantities are written as $v_c \propto t^{-1/2}$, $v_h \propto \text{const.}$, $P_c \propto t^{-1}$, $A_h \propto \text{const.}$ The absolute values of each physical quantities are also agree with those in BC89 when omitting $1/(\hat{\gamma}_c - 1)$ and replace $C_{vc} = 1/\pi^{0.5}$, $C_{vh} = 1$, $C_{pc} = 1$, and $C_{ah} = 1$. It is included in Fig. 2 the case of BC89 with the parameters of $\alpha = 0$, $X = 1$, and $A_h = 30 \text{ kpc}^2$. As explained before, the larger (smaller) α predict smaller (larger) L_j because of the dilute (dense) ICM. The comparison with the analytic model and numerical studies, is also important issue. By Cioffi and Blondin (1992), the issues of “head cross section growth” and “decreasing head velocity” is assessed by hydrodynamic simulation and their result shows $A_h \propto t^{0.4}$. As a good example, our solution of $X \simeq 1.1$ describes the studies Nath (1995) corresponds to case of flat solution above $v_h \propto t^{-0.2}$, $v_c \propto t^{-0.45}$, $P_c \propto t^{-0.9}$, and $A_h \propto t^{0.4}$. Concerning A_c , since most of numerical studies have mainly focused on the propagation of cylindrical geometry jets (e.g., Marti et al. 1997; Clarke, Harris and Carilli 1997), it remains as an important future work on the time evolution of A_c .

3 TOTAL KINETIC POWER AND SOURCE AGE OF CYGNUS A

Here we explore L_j and t_{age} by matching the observed cross-sectional areas and lengths of a cocoon (i.e., r_h , r_c , A_h , and A_c). In this paper, we focus on the archetypal radio galaxy Cygnus A (Carilli and Barthel 1996; Carilli and Harris 1996 for reviews).

First, we estimate the typical values of physical quantities. Concerning the mass density profile of ICM, we adopt $\alpha = 1.5$ based on Reynolds and Fabian (1996) and Smith et al. (2002). We examine the case of $X = 12/7$ as a fiducial one which predict the constant $\mathcal{R}$ in time. Other observed quantities $\bar{\rho}_a = 0.5 \times 10^{-2} m_p \text{ g cm}^{-3}$, and $r_h = 60 \text{ kpc}$, based on Carilli et al. (1998), Arnaud et al. (1984), and Smith et al. (2002). Here we assume $\hat{\gamma}_c = 4/3$. Using these quantities, we obtain

$$\beta_h(t) = 8.36 \times 10^{-3} L_{j,46} t_{20}^2 \left(\frac{t}{t_{20}} \right)^{-1/7}, \quad (9)$$

$$\beta_c(t) = 6.84 \times 10^{-3} t_{20}^{-1} \left(\frac{t}{t_{20}} \right)^{-1/7}, \quad (10)$$

$$P_c(t) = 4.78 \times 10^{-10} t_{20}^{-2} \left(\frac{t}{t_{20}} \right)^{-11/7} \text{ dyne cm}^{-2}, \quad (11)$$

$$A_h(t) = 66.4 L_{j,46}^{1/2} t_{20}^{1/2} \left(\frac{t}{t_{20}} \right)^{11/7} \text{ kpc}^2, \quad (12)$$

where $\beta_h = v_h/c$, $\beta_c = v_c/c$, $t_{20} = t_{\text{age}}/20 \text{ Myr}$, $L_{j,46} = L_j/10^{46} \text{ erg s}^{-1}$, with the resultant value of $\mathcal{R}(t_{\text{age}}) = 0.815$ for Cygnus A. Note that $\mathcal{R} \propto \bar{\beta}_h/\bar{\beta}_c$ has a strong dependence on the age such as $\propto t_{\text{age}}^3$, the allowed source age is fairly restricted.

Next, to clarify the allowed range of t_{age} and L_j , we impose the following conditions; the condition which should be satisfied is that (I) $\mathcal{R}(t_{\text{age}}) \sim 0.5 - 0.7$, from the *Chandra* image (Wilson et al. 2000), (II) cocoon pressure is over-pressured $P_c > P_a = 8 \times 10^{-11} \text{ dyn cm}^{-2}$ (Arnaud et al. 1984; Smith et al. 2002), (III) the area size of A_h lies in the range of $30 \text{ kpc}^2 < A_h < 150 \text{ kpc}^2$. The minimum value corresponds to the one adopted in BC89. From the radio image of Perley, Dreher and Cowan (1984), we employ the maximum value as $A_h = 150 \text{ kpc}^2$, which corresponds to the cross sectional area of the radio lobe at the location of the hot spot. In Fig. 4, we show the resultant source age and total kinetic power of the jet. The region of the source age larger than $\sim 30 \text{ Myr}$ is not allowed by the condition (II). Larger (smaller) A_h predict larger (smaller) L_j and younger t_{age} . Obtained values are $2 \times 10^{45} \text{ erg s}^{-1} < L_j < 1 \times 10^{48} \text{ erg s}^{-1}$ and $3 \text{ Myr} < t_{\text{age}} < 30 \text{ Myr}$.

Let us consider how the uncertainties of L_j and t_{age} are determined. From Eqs. (7) and (8), $A_h \propto L_j t_{\text{age}}^2$ and $\mathcal{R} \propto L_j^{-1/(\alpha-4)} t_{\text{age}}^{-3/(\alpha-4)}$ are obtained. Since A_h and $\mathcal{R}$ have uncertainties with the factors of 5 and 1.4 respectively, the allowed region is mainly controlled by A_h . In the present work, in view of the comparison of BC89, we took the minimum value as $A_h = 30 \text{ kpc}^2$. However, from the hydrodynamical point of view, it seems natural to suppose that A_h to be the close value to the cross sectional area of the radio lobe at the distance of head, i.e., $A_h = 150 \text{ kpc}^2$.

3.1 Comparison with previous works

The resultant age well agree with the independent result of synchrotron age model by Carilli et al. (1991), which claims that $6 \text{ Myr} < t_{\text{age}} < 30 \text{ Myr}$. The velocity of the hot spot $\beta_{\text{hs}} \sim 0.06$ corresponds to the source age of 6 Myr, while $\beta_{\text{hs}} \sim 0.01$ corresponds to the source age of 30 Myr.

As a complementary result, L_j estimated in KT04 in the range of $6 \text{ Myr} < t_{\text{age}} < 30 \text{ Myr}$ based on the result Carilli et al. (1991) is shown in Fig. 4 (the solid line). KT04 estimate the total kinetic power of the relativistic jet as

$$\begin{aligned} L_j &= A_j c \Gamma_j^2 \beta_j \rho_j c^2 \\ &= A_j c \left(\frac{r_{60}}{t_{\text{age}}} \right)^2 \rho_a(r_{60}) \propto A_j \end{aligned}$$

in the strong relativistic shock limit, where $A_j = \pi R_{\text{hs}}^2$, Γ_j , $\beta_j c$, ρ_j , and $R_{\text{hs}} = 2 \text{ kpc}$ (Wilson et al. 2000) are the cross-sectional area of the jet the Lorentz factor, the velocity, and mass density of the jet, and the hot spot radius, respectively. It should be stressed that the cocoon model can predict both L_j and t_{age} at the same time, while the synchrotron aging model (Carilli et al. 1991) and the 1D jet model (KT04) only

determine t_{age} or L_j . Although these three models are independent, they show a reasonable agreement on the values of L_j and t_{age} with each other at least in order-of-magnitude in the case of Cygnus A. At the same time, it is worth to discuss a factor of deviation of the estimate L_j by the cocoon model and 1D jet model. In the same way as KT04, 1D analysis between the jet and ICM take the plane parallel assumption (e.g., Begelman, Blandford & Rees 1984). The approximation means that A_j equals to A_h . However, A_h is supposed to be larger than A_j . Hence, the plane parallel approximation would cause the underestimate of the L_j in KT04.

Rawlings & Saunders (1991) (hereafter RS91) is the pioneering work on the correlation between the L_j and the luminosity of the narrow line regions, which lies in close to the central engine. Then the comparison with the present work and RS91 is intriguing issue. For the sources where synchrotron spectral aging are available, it is simply by

$$Q = \frac{E_{\text{eq}}}{t_{\text{age}}\eta}, \quad \eta = 0.5$$

where Q , η , and E_{eq} are the kinetic power of jet, a parameter expressing the fraction of the work done on the ICM, and the equipartition energy with the electrons and the magnetic field make an equal contribution to the total energy density (e.g., Miley (1980)), respectively. We focus on the sample sources where t_{age} is independently obtained by synchrotron aging method. One difference between the present work and RS91 is that we solved the equations of motion and energy equation (3 Eqs. in total), while the RS91 only employ energy equation. Because of this, we can eliminate the free parameter η . More important and essential difference is that Q in RS91 is not a total kinetic power but it is just a equipartition power (even though they insist that it as a total power). We emphasize the advantage of our work of dealing with the total kinetic power whilst RS91 only handles the equipartition power of extragalactic jets.

Kaiser (2000) (hereafter K00) independently addressed this quantity by studying some bright radio sources involving Cygnus A. Compared with RS91, the common advantage is that both K00 and the present work can estimate the total kinetic power by directly dealing with the hydrodynamics. The way of comparing the observation with the model by K00 is different from ours. K00 matched the surface brightness distribution of the cocoon along the jet axis. The main difference between the present work and K00 is the derived cocoon pressure during the matching of the observations and the models. In the example of Cygnus A, compared with our estimate of $P_c \sim 5 \times 10^{-10} \text{ erg cm}^{-3}$, they tend to estimate smaller P_c of order of $P_c \sim 10^{-12} - 10^{-11} \text{ erg cm}^{-3}$ (Tables 2, 3 and 4 in K00). This mainly cause the deviation of the derived total kinetic power of Cygnus A as $L_j \sim (\text{a few}) \times 10^{46} \text{ erg s}^{-1}$, whilst K00 derive the total kinetic power with order of $L_j \sim (\text{a few}) \times 10^{45} \text{ erg s}^{-1}$.

3.2 Efficiency of accretion power to kinetic power

In Tadhunter et al. (2003), the mass of the super-massive black hole (SMBH) of Cygnus A is reported as $2.5 \times 10^9 M_\odot$ which leads to the Eddington luminosity as $L_{\text{Edd}} = 3 \times 10^{47} \text{ erg s}^{-1}$. From this it follows that $L_j/L_{\text{Edd}} \sim 0.01 - 1$. On the basis of the observational evidence for the accretion-flow ori-

gin for the jet in AGNs (e.g., Marscher et al. 2002), it is clear that the value of L_j/L_{Edd} directly shows the required minimum rate of the mass accretion onto the SMBH normalized by the corresponding Eddington mass accretion rate.

Merloni, Heinz and Di Matteo 2003 (see also Maccarone, Gallo and Fender 2003) examine the disk-jet connection by studying the correlation between the radio (L_R) and the X-ray (L_X) luminosity and the black hole mass. With the large samples of the stellar mass black holes and SMBHs, they claim these sources have the “fundamental plane” in the 3 dimensional space of these quantities. On the way, the quantity of L_X/L_{Edd} is brought up to probe the activity of the central engine. For Cygnus A, Young et al. (2002) reports $L_X \simeq 3.7 \times 10^{44} \text{ erg s}^{-1}$ which leads to $L_X/L_{\text{Edd}} \sim 10^{-3}$. Whereas we recognize the usability of the quantity of L_X/L_{Edd} to probe the engine activity, we emphasize that the L_X largely depends on the accretion flow model and the radiation efficiency at X-ray band. On the other hand, the quantity of L_j/L_{Edd} addressed in the present work does not have the model dependence and directly shows us the engine activity.

Maraschi & Tavecchio (2003) also addressed this quantity by collecting large samples of blazars. They estimated the jet power from the blazar’s non-thermal spectrum energy distribution, which are well understood as synchrotron plus inverse Compton (e.g., Maraschi et al. 1992; Sikora et al. 1994; Blandford & Levinson 1995; Kino et al. 2002). The upper limit of the accretion power is inferred from the luminosity of observed broad emission lines. They conclude that for flat-spectrum radio quasars (FSRQs) the the total power of the jet is of the same order as the accretion power. Note that their estimate of the accretion power from the emission line is putative way and the estimated total power does not include the contribution of protons associated with thermal electrons which may cause some underestimate of the total power of the jet.

We again emphasize that as we see above, the value of L_j/L_{Edd} for Cygnus A in the present work is obtained with fewer assumptions. Moreover it make us a fairly robust probe of the central engine activity of AGN jets. The application of our method to a large sample of other AGN jets surely bring about new and important knowledges. This is actually our ongoing project.

4 CONCLUSION

Our main conclusions in the present work are as follows;

(i) A new method to estimate the total kinetic power of the jet L_j and source age t_{age} in powerful FR II radio sources is proposed. For that, the study of cocoon dynamics by BC89 is revisited and physical quantities associated with the cocoon are analytically solved as functions of L_j and t_{age} . The comparison of the analytic solution with the observed cocoon shape lead to the L_j and t_{age} in general.

(ii) The analysis is focused on Cygnus A, with the conditions of $0.5 \leq \mathcal{R} \leq 0.7$ and $30 \text{ kpc}^2 < A_h < 150 \text{ kpc}^2$. The estimated age $3 \text{ Myr} < t_{\text{age}} < 30 \text{ Myr}$ shows a good agreement of the independently estimate age by synchrotron aging model by Carilli et al. (1991). The estimated total kinetic power $0.2 \times 10^{46} \text{ erg s}^{-1} < L_j < 1 \times 10^{48} \text{ erg s}^{-1}$ has

also a reasonable agreement with the independent 1D jet model of KT04 with the aid of $6 \text{ Myr} < t_{\text{age}} < 30 \text{ Myr}$.

(iii) For Cygnus A, we find that the total kinetic power lies in the range of $L_j/L_{\text{Edd}} \sim 0.01 - 1$, while the X -ray luminosity $L_X \simeq 3.7 \times 10^{44} \text{ erg s}^{-1}$ (Young et al. 2002) satisfies $L_X/L_{\text{Edd}} \sim 10^{-3}$. The result of $L_j/L_{\text{Edd}} \sim 0.01 - 1$ indicate the lower limit of the mass accretion rate, which gives the crucial hint for resolving the jet formation problem.

ACKNOWLEDGMENTS

We appreciate the helpful comments of the referee to improve the paper. We thank A. C. Fabian, D. E. Harris, and D. Schwartz for valuable comments. M. K. thank A. Celotti, J. Kataoka, N. Isobe and A. Mizuta for stimulating discussions. We acknowledge the Italian MIUR and INAF financial supports.

REFERENCES

- Arnaud K. A., Fabian A. C., Eales S. A., Jones C., Forman W., 1984, MNRAS, 211, 981
- Begelman, M. C., Blandford, R. D., & Rees, M. J. 1984, Rev. Mod. Phys., 56, 255
- Begelman, M. C. & Cioffi, D. F. 1989, ApJL, 345, L21 (BC89)
- Bicknell G. V., Dopita M. A., O'Dea C. P. O., 1997, ApJ, 485, 112
- Blandford R. D., Levinson A., 1995, ApJ, 441, 79
- Carilli, C. L., Perley, R. A., Dreher, J. W., & Leahy, J. P. 1991, ApJ, 383, 554
- Carilli, C. L. & Barthel, P. D. 1996, A&ARv, 7, 1
- Carilli C. L., & Harris D. E., 1996, Proc. Greenbank Workshop, Cygnus A - Study of Radio Galaxy. Cambridge Univ. Press, Cambridge
- Carilli C. L., Perley R., Harris D. E., Barthel P. D., 1998, Physics of Plasmas, 5, 1981
- Celotti A., Ghisellini G., Chiaberge M., 2001, MNRAS, 321, L1
- Celotti A., Kuncic Z., Rees M. J., Wardle J. F. C., 1998, MNRAS, 293, 288
- Cioffi D. F., Blondin J. M., 1992, ApJ, 392, 458
- Clarke D. A., Harris D. E., Carilli C. L., 1997, MNRAS, 284, 981
- Fanti C., Fanti R., Dallacasa D., Schilizzi R. T., Spencer R. E., Stanghellini C., 1995, A&A, 302, 317
- Croston, J. H., Hardcastle, M. J., Harris, D. E., Belsole, E., Birkinshaw, M., Worrall, D. M., 2005, ApJ, 626, 733
- Dunn R. J. H., Fabian A. C., 2004, MNRAS, 355, 862
- Fabian A. C., Celotti A., Blundell K. M., Kassim N. E., Perley R. A., 2002, MNRAS, 331, 369
- Hardcastle M. J., Birkinshaw M., Cameron R. A., Harris D. E., Looney L. W., Worrall D. M., 2002, ApJ, 581, 948
- Harris D. E., Krawczynski H., 2002, ApJ, 565, 244
- Isobe N., Tashiro M., Makishima K., Iyomoto N., Suzuki M., Murakami M. M., Mori M., Abe K., 2002, ApJ, 580, L111
- Kaiser C. R., 2000, A&A, 362, 447
- Kaiser C. R., Alexander P., 1997, MNRAS, 286, 215
- Kataoka J., Leahy J. P., Edwards P. G., Kino M., Takahara F., Serino Y., Kawai N., Martel A. R., 2003, A&A, 410, 833
- Kino M., Takahara F., Kusunose M., 2002, ApJ, 564, 97
- Kino, M. & Takahara, F. 2004, MNRAS, 349, 336 (KT04)
- Krause M., 2003, A&A, 398, 113
- Leahy J. P., Gizani N. A. B., 2001, ApJ, 555, 709
- Nath B. B., 1995, MNRAS, 274, 208
- Maccarone T. J., Gallo E., Fender R., 2003, MNRAS, 345, L19




Figure 1. A cartoon of interaction of the ICM with declining atmosphere and the relativistic jet in FR II radio galaxy. As a result, most of the kinetic energy of jet is deposited in the cocoon and it is inflated by its internal energy.

- Marscher A. P., Jorstad S. G., Gómez J., Aller M. F., Teräsranta H., Lister M. L., Stirling A. M., 2002, Nature, 417, 625
- Maraschi L., Ghisellini G., Celotti A., 1992, ApJ, 397, L5
- Maraschi L., Tavecchio F., 2003, ApJ, 593, 667
- Marti J. M. A., Mueller E., Font J. A., Ibanez J. M. A., Marquina A., 1997, ApJ, 479, 151
- Merloni A., Heinz S., di Matteo T., 2003, MNRAS, 345, 1057
- Miley G., 1980, ARA&A, 18, 165
- Perley R. A., Dreher J. W., Cowan J. J., 1984, ApJ, 285, L35
- Rawlings S., Saunders R., 1991, Nature, 349, 138
- Readhead A. C. S., Taylor G. B., Xu W., Pearson T. J., Wilkinson P. N., Polatidis A. G., 1996a, ApJ, 460, 612
- Readhead A. C. S., Taylor G. B., Pearson T. J., Wilkinson P. N., 1996b, ApJ, 460, 634
- Reynolds C. S., Fabian A. C., 1996b, MNRAS, 278, 479
- Scheuer P. A. G., 1974, MNRAS, 166, 513
- Sikora M., Begelman M. C., Rees M. J., 1994, ApJ, 421, 153
- Sikora M., Madejski G., 2000, ApJ, 534, 109
- Smith D. A., Wilson A. S., Arnaud K. A., Terashima Y., Young A. J., 2002, ApJ, 565, 195
- Tadhunter C., Marconi A., Axon D., Wills K., Robinson T. G., Jackson N., 2003, MNRAS, 342, 861
- Tavecchio F., Maraschi L., Sambruna R., Urry C. M., 2000, ApJ, 544, L23
- Wilson A. S., Young A. J., Shopbell P. L., 2000, ApJ, 544, L27
- Young A. J., Wilson A. S., Terashima Y., Arnaud K. A., Smith D. A., 2002, ApJ, 564, 176




Figure 2. The allowed regions of L_j and t_{age} with the different values of α with $X = 1$. The region is determined by the constraints of $0.5 < \mathcal{R} < 1$ and $30\text{kpc}^2 < A_h < 150\text{kpc}^2$. We examine $\alpha = 0$ and 2 here. The case of $\alpha = 0$, and $A_h = 30\text{kpc}^2$ by Begelman and Cioffi (1989) is involved in the right-lower part of the filled region. Larger α requires significantly larger L_j for plowing the larger amount of ICM.




Figure 3. The allowed regions of L_j and t_{age} with the different values of X with $\alpha = 1.5$. Likewise Fig. 2, we adopt $0.5 < \mathcal{R} < 1$ and $30\text{kpc}^2 < A_h < 150\text{kpc}^2$. The case of $X = 1.62$, $12/7$ and 1.80 are shown here. Larger X requires smaller L_j corresponding to the slower velocity of the sideways expand.




Figure 4. The allowed region of L_j and t_{age} of Cygnus A (filled in black). The under-pressured region $P_c < P_a$ is excluded by definition. The case of $0.5 < \mathcal{R} < 0.7$ and $30\text{kpc}^2 < A_h < 150\text{kpc}^2$ is examined. As a reference, Eddington luminosity and the total kinetic power of jet estimated in KT04 are shown in the thick-solid and solid lines, respectively.